

On Existential Observability and Reachability in a Class of Discrete-Time Switched Linear Systems

Magnus Egerstedt and Mohamed Babaali

Abstract—We study the class of switched linear systems with constant dynamics, i.e. with switching in only the measurement or control matrices (which corresponds to systems with several sensory and actuation modes), and establish that the length of the shortest mode path achieving observability or reachability, if one exists, is at most the square of the dimension of the state space. This not only proves that observability/reachability is decidable for such systems, but it also gives a direct way of finding the observable/reachable mode path.

I. RESULT

We are concerned, in this paper, with systems of the form

$$\begin{aligned} x_{k+1} &= Ax_k + B(\theta_k)u_k \\ y_k &= C(\theta_k)x_k \end{aligned} \quad (1)$$

where x_k , y_k and u_k are in \mathbb{R}^n , \mathbb{R}^p and \mathbb{R}^m , respectively, where A , $C(\cdot)$ and $B(\cdot)$ are real matrices of compatible dimensions, and where the discrete mode $\theta_k \in \bar{s} \triangleq \{1, \dots, s\}$, so that $B(\theta_k) \in \{B(1), \dots, B(s)\}$ and $C(\theta_k) \in \{C(1), \dots, C(s)\}$, which are fixed sets of matrices. We ask whether there exists a *path*, i.e., a mode string over \bar{s} , $\theta \triangleq \theta_1 \dots \theta_N$, such that the matrix

$$C(\theta) \triangleq (B(\theta_N) B(\theta_{N-1})A \dots B(\theta_1)A^{N-1}) \quad (2)$$

has full (row) rank. We refer to such paths as *reachable paths*, since the reachable set of (1) under such paths is the whole state space. By duality, we end up also considering the equivalent problem of asking whether there exists an *observable path*, i.e. a path $\theta = \theta_1 \dots \theta_N$ such that

$$\mathcal{O}(\theta) \triangleq \begin{pmatrix} C(\theta_1) \\ \vdots \\ C(\theta_N)A^{N-1} \end{pmatrix} \quad (3)$$

has full (column) rank, making it possible to observe the initial state x_1 in (1) under the path θ . We will refer to the length of the minimum-length observable (resp. reachable) path of some system as its *index* of observability (resp. reachability). The purpose of this paper is to establish the following result.

Main Result. *If (1) is observable (resp. reachable) then its index of observability (resp. reachability) is at most n^2 . \diamond*

Magnus Egerstedt's work was sponsored by NSF through the ECS NSF-CAREER award (grant # 0237971). The research of Mohamed Babaali was partially supported by NSF CAREER Grant 0132716.

Magnus Egerstedt is with the School of Electrical and Computer Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA (magnus@ece.gatech.edu). Mohamed Babaali is with the School of Engineering and Applied Science, University of Pennsylvania, Philadelphia, PA 19104, USA (babaali@grasp.cis.upenn.edu).

This result clearly implies that reachability (resp. observability) is decidable in our class of switched linear systems, since it suffices to check whether there exists a reachable (resp. observable) path of length n^2 . After giving a short overview of the background to this problem in Section II, we prove our Main Result in Section III.

II. BACKGROUND

We first note that most of the existing literature is focused on the reachability problem, and we will report it as such. It should also be stressed that our systems (1) form a special class of switched linear systems

$$x_{k+1} = A(\theta_k)x_k + B(\theta_k)u_k, \quad (4)$$

where the A matrix also undergoes switching, so that $A(\theta_k) \in \{A(1), \dots, A(s)\}$.

The problem of determining the existence of a reachable path for (4) has eluded researchers for a long time, and, to the best of our knowledge, the only efforts in studying the existence of reachable paths for (4) can be found in [2], [3], [4], [5], [6]. First, in a series of papers [2], [3], [6], Stanford and Conner studied the structure of the reachable set of switched linear systems, and showed that it was a subspace of \mathbb{R}^n in *reversible* (i.e. with invertible $A(\theta_k)$ matrices) systems [6], and, in general, a union of arbitrarily many maximal subspaces in [3]. They also, in parallel, tried to find upper bounds on the index of reachability depending only on s , n and m . Unfortunately, they only obtained partial results for special classes, i.e., for special combinations of n , m and s and for classes of switched linear systems with rank constraints on the parameters $B(i)$ and $A(i)$. In short, it is still unknown whether the indexes of reachability are bounded, even for the class reversible systems. However, the existence of a reachable path in reversible systems has recently been shown to be decidable in [4], [5], although not constructively (no upper bounds on the indexes were given). More precisely, it was shown that a reversible system was reachable if and only if the minimal subspace invariant with respect to every $A(i)$, $1 \leq i \leq s$, and containing $\text{Im}(B(1)) + \dots + \text{Im}(B(s))$, equals the whole state space \mathbb{R}^n . This problem is clearly decidable since the condition above can easily be cast into a finite matrix rank problem (See [4]). Therefore, since A in (1) is not necessarily invertible, this paper is the first to establish the decidability of reachability for our class, and moreover, it is the first to give upper bounds for such a broad class of systems. Note, however, that the general problem remains open.

It is beyond the scope of this paper to provide an exhaustive literature survey on observability and controllability in switched and hybrid systems, which has become plethoric. However, closely related problems include the continuous-time version of the problem we consider in this paper, which was shown to be decidable in [5], [7], and the problem of determining the existence of an integer N such that every path of length N is reachable, which is the “universal” version of the “existential” problem considered here. This problem has been shown to be decidable, independently, in [1] and [5], where, furthermore, the indexes of *pathwise controllability* were shown to be bounded by numbers depending only on s and n .

III. PROOF OF MAIN RESULT

We will prove the observability version of the result, since it makes the presentation more straightforward. We first state, without proof, the following immediate corollary to the Cayley-Hamilton Theorem.

Lemma 1. *If $A \in \mathbb{R}^{n \times n}$, then whenever $i \geq n$, there exists a set of real numbers $\{\alpha_i\}_{j=1}^n$ such that $A^i = \sum_{j=1}^n \alpha_j A^{j-1}$.* \diamond

Next, in order to ease the discussion, we continue with with a few definitions. Let $\mathcal{R}(M)$ denote the row range space of a matrix M and $\rho(M)$ its rank. $|\theta|$ denotes the length of θ . θ^1 and θ^2 being paths of length N_1 and N_2 respectively, $\theta^1\theta^2$ denotes their concatenation $\theta_1^1\theta_2^1 \cdots \theta_1^2\theta_2^2 \cdots \theta_{N_2}^2$. Furthermore, $\theta^{(q)}$ is the path θ concatenated with itself $q-1$ times. Given a path θ , $\theta_{[i,j]}$ is its substring (or infix) $\theta_i\theta_{i+1} \cdots \theta_j$. By convention, we let $\theta_{[i,i-1]} = \epsilon$, the null string, for all $1 \leq i \leq |\theta|$. Finally, let $\mathcal{O}(\theta)_{[i,j]}$ denote the submatrix of $\mathcal{O}(\theta)$ constituted by rows i through j of $\mathcal{O}(\theta)$:

$$\mathcal{O}(\theta)_{[i,j]} \triangleq \begin{pmatrix} C(\theta_i)A^{i-1} \\ \vdots \\ C(\theta_j)A^{j-1} \end{pmatrix},$$

and note that $\mathcal{O}(\theta)_{[i,j]} = \mathcal{O}(\theta_{[i,j]})A^{i-1}$, for all i and j such that $1 \leq i \leq j \leq |\theta|$. Let $\mu_k(\theta) \triangleq \rho(\mathcal{O}(\theta_{[1,k]}))$, and note that $k \mapsto \mu_k(\theta)$ is nondecreasing with at most n jumps. Let $N(\theta)$ be the number of jumps of $\mu_k(\theta)$, and let $i_l(\theta)$ be the time of the l^{th} jump of $\mu_k(\theta)$. To simplify matters in the sequel, we let $i_0(\theta) \triangleq 0$. Finally, let $I(\theta) \triangleq \{l \mid i_{l+1}(\theta) - i_l(\theta) > n\}$ index the set of consecutive jumps spaced by more than n time instants. If $I(\theta) \neq \emptyset$, let $q(\theta) \triangleq \min I(\theta)$ denote the position (not the time) of the first jump preceding the jump following it by more than n time units. Note that $I(\theta) \subset \{1, \dots, n\}$ and $q(\theta) \leq n$ for any θ , since $\mu_k(\theta)$ has at most n jumps.

Lemma 2. *If θ is observable and $I(\theta) \neq \emptyset$, then, letting $l = q(\theta)$,*

$$\theta' = \theta_{[1, i_l(\theta)]} \theta_{[i_{l+1}(\theta)]}^{(n)} \theta_{[i_l(\theta)+n+1, |\theta|]} \quad (5)$$

is observable with either $I(\theta') = \emptyset$ or $q(\theta') > q(\theta)$. \diamond

Proof: First, $l \in I(\theta)$ implies that $i_{l+1}(\theta) - i_l(\theta) > n$, which, by Lemma 1, gives real numbers $\{\alpha_i\}_{j=1}^n$ such that $A^{i_{l+1}(\theta)-i_l(\theta)-1} = \sum_{j=1}^n \alpha_j A^{j-1}$, thus that $C(\theta_{i_{l+1}(\theta)})A^{i_{l+1}(\theta)-i_l(\theta)-1} = \sum_{j=1}^n \alpha_j C(\theta_{i_{l+1}(\theta)})A^{j-1}$, which, right-multiplied by $A^{i_l(\theta)}$ on both sides, gives $C(\theta_{i_{l+1}(\theta)})A^{i_{l+1}(\theta)-1} = \sum_{j=1}^n \alpha_j C(\theta_{i_{l+1}(\theta)})A^{i_l(\theta)+j-1}$, which is equivalent to $\mathcal{R}(C(\theta_{i_{l+1}(\theta)})A^{i_{l+1}(\theta)-1}) \subset \mathcal{R}(\mathcal{O}(\theta')_{[i_l(\theta)+1, i_l(\theta)+n]})$. By definition of θ' , we furthermore have $\mathcal{R}(\mathcal{O}(\theta')_{[1, i_l(\theta)]}) = \mathcal{R}(\mathcal{O}(\theta)_{[1, i_l(\theta)]})$. Moreover, $\mu_k(\theta)$ having a jump at $i_{l+1}(\theta)$ implies that $\mathcal{R}(C(\theta_{i_{l+1}(\theta)})A^{i_{l+1}(\theta)-1}) \not\subset \mathcal{R}(\mathcal{O}(\theta)_{[1, i_l(\theta)]})$. Combining the last three claims, we get $\mathcal{R}(\mathcal{O}(\theta')_{[i_l(\theta)+1, i_l(\theta)+n]}) \not\subset \mathcal{R}(\mathcal{O}(\theta')_{[1, i_l(\theta)]})$, which, since $i_j(\theta') = i_j(\theta)$ for $j \leq l$ because $\theta'_{[1, i_l(\theta)]} = \theta_{[1, i_l(\theta)]}$, implies that $i_{l+1}(\theta') - i_l(\theta') \leq n$, proving that either $I(\theta') = \emptyset$ or $q(\theta') > l = q(\theta)$.

It now remains to show that $\mathcal{R}(\mathcal{O}(\theta)) \subset \mathcal{R}(\mathcal{O}(\theta'))$, which, since θ is observable, resulting in $\rho(\mathcal{O}(\theta)) = n$, would imply that $\rho(\mathcal{O}(\theta')) = n$, hence that θ' is observable as well. To this end, we first note that, by definition of θ' , $\theta_{[1, i_l(\theta)]} = \theta'_{[1, i_l(\theta)]}$ and $\theta_{[i_l(\theta)+n+1, |\theta|]} = \theta'_{[i_l(\theta)+n+1, |\theta'|]}$, which implies that $\mathcal{R}(\mathcal{O}(\theta)_{[1, i_l(\theta)]}) = \mathcal{R}(\mathcal{O}(\theta')_{[1, i_l(\theta)]}) \subset \mathcal{R}(\mathcal{O}(\theta'))$ and that $\mathcal{R}(\mathcal{O}(\theta)_{[i_l(\theta)+n+1, |\theta|]}) = \mathcal{R}(\mathcal{O}(\theta')_{[i_l(\theta)+n+1, |\theta'|]}) \subset \mathcal{R}(\mathcal{O}(\theta'))$. Therefore, all that remains to be shown is that $\mathcal{R}(\mathcal{O}(\theta)_{[i_l(\theta)+1, i_l(\theta)+n]}) \subset \mathcal{R}(\mathcal{O}(\theta'))$, which follows from $\mathcal{R}(\mathcal{O}(\theta)_{[i_l(\theta)+1, i_l(\theta)+n]}) \subset \mathcal{R}(\mathcal{O}(\theta)_{[1, i_l(\theta)]})$, (which, itself, is due to $i_{l+1}(\theta) - i_l(\theta) > n$) combined with $\mathcal{R}(\mathcal{O}(\theta)_{[1, i_l(\theta)]}) \subset \mathcal{R}(\mathcal{O}(\theta'))$. \square

Proof of Main Result: Let θ be observable. If $I(\theta) = \emptyset$, then we are done, since $i_{N(\theta)} \leq n^2$, and therefore $\theta_{[1, n^2]}$ is observable of length n^2 . Otherwise, let θ' be given by lemma 2. Then θ' is observable, and furthermore, either $I(\theta')$ is empty, in which case we are done, or $q(\theta') > q(\theta)$, in which case we can, once again, invoke Lemma 2 and get an observable θ'' such that $q(\theta'') > q(\theta') > q(\theta)$ or $I(\theta'') = \emptyset$. Finally, since $q(\cdot)$ is bounded by n , the previous procedure will eventually end, yielding an observable path λ such that $I(\lambda) = \emptyset$, thus that $\lambda_{[1, n^2]}$ is observable. \square

REFERENCES

- [1] M. Babaali and M. Egerstedt, “Pathwise observability and controllability are decidable,” in *Proceedings of the 42nd IEEE Conference on Decision and Control*, Maui, HI, December 2003, pp. 5771–5776.
- [2] L. T. Conner, Jr. and D. P. Stanford, “State deadbeat response and observability in multimodal systems,” *SIAM Journal on Control and Optimization*, vol. 22, no. 4, pp. 630–644, 1984.
- [3] —, “The structure of the controllable set for multimodal systems,” *Linear Algebra and its Applications*, vol. 95, pp. 171–180, 1987.
- [4] S. S. Ge, Z. Sun, and T. H. Lee, “Reachability and controllability of switched linear discrete-time systems,” *IEEE Transactions on Automatic Control*, vol. 46, pp. 1437–1441, September 2001.
- [5] L. Gurvits, “Stabilities and controllabilities of switched systems (with applications to the quantum systems),” in *Proceedings of the Fifteenth International Symposium on Mathematical Theory of Networks and Systems*, Univ. Notre Dame, August 2002.
- [6] D. P. Stanford and L. T. Conner, Jr., “Controllability and stabilizability in multipair systems,” *SIAM Journal on Control and Optimization*, vol. 18, no. 5, pp. 488–497, 1980.
- [7] Z. Sun, S. S. Ge, and T. H. Lee, “Controllability and reachability criteria for switched linear systems,” *Automatica*, pp. 775–786, May 2002.